




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C09B 1/515	A1	(11) International Publication Number: WO 91/05824 (43) International Publication Date: 2 May 1991 (02.05.91)
(21) International Application Number: PCT/GB90/01574 (22) International Filing Date: 12 October 1990 (12.10.90) (30) Priority data: 8923075.9 13 October 1989 (13.10.89) GB (71) Applicant (for all designated States except US): NATIONAL RESEARCH DEVELOPMENT CORPORATION [GB/GB]; 101 Newington Causeway, London SE1 6BU (GB). (72) Inventor; and (75) Inventor/Applicant (for US only) : PATTERSON, Laurence, Hylton [GB/GB]; 44 Glenfield Frith Drive, Glenfield, Leicester LE3 8PQ (GB). (74) Agent: STEPHENSON, Gerald, Frederick; Patents Department, National Research Development Corporation, 101 Newington Causeway, London SE1 6BU (GB).		(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), US. Published With international search report.

(54) Title: ANTI-CANCER COMPOUNDS



(I)

(57) Abstract

Compounds of formula (I) in which R₁, R₂, R₃ and R₄ are each separately selected from hydrogen, X, NH-A-NHR and NH-A-N(O)R'R'' wherein X is hydroxy, halogeno, amino, C₁₋₄ alkoxy or C₂₋₈ alkanoyloxy, A is a C₂₋₄ alkylene group with a chain length between NH and NHR or N(O)R'R'' of a least 2 carbon atoms and R, R' and R'' are each separately selected from C₁₋₄ alkyl groups and C₂₋₄ hydroxyalkyl and C₂₋₄ dihydroxyalkyl groups in which the carbon atom attached to the nitrogen atom does not carry a hydroxy group and no carbon atom is substituted by two hydroxy groups, or R' and R'' together are a C₂₋₆ alkylene group which with the nitrogen atom to which R' and R'' are attached forms a heterocyclic group having 3 to 7 atoms in the ring, but with the proviso that at least one of R₁ to R₄ is a group NH-A-N(O)R'R'', and physiologically acceptable salts thereof are of value in the treatment of cancer.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SN	Senegal
CH	Switzerland	LI	Liechtenstein	SU	Soviet Union
CI	Côte d'Ivoire	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

- 1 -

ANTI-CANCER COMPOUNDS

This invention relates to novel anthraquinones which are of particular value in the treatment of cancer.

05 A wide variety of aminoalkylamino anthraquinones (aminoalkyl-aminoanthracene-9,10-diones) has been described for use as chemotherapeutic agents for the treatment of cancer, perhaps the most active being the compound mitoxantrone (mitozantrone) of formula




which is the subject of U.S. Patent 4,197,249 and U.K. Patent 2,004,293B. However, in common with other cytotoxic
10 chemotherapeutic agents the aminoalkylamino anthraquinones have the disadvantage that their activity is not confined to neoplastic cells and they therefore exhibit various undesirable side effects including, to a greater or lesser extent among the different compounds, myelosuppression and cardiotoxicity.

15 It is an object of the present invention to provide a group of anthraquinone pro-drugs which are of lesser cytotoxicity than the drug itself, preferably being substantially non-cytotoxic, the pro-drugs being converted in vivo under the anaerobic conditions within neoplastic tissue to the cytotoxic drug thereby mitigating
20 the side effects of administering that drug directly.

- 2 -

Accordingly the present invention comprises a compound of formula (I)



(I)

in which R_1 , R_2 , R_3 and R_4 are each separately selected from hydrogen, X, $NH-A-NHR$ and $NH-A-N(O)R'R''$ wherein X is hydroxy, halogeno, amino, C_1-4 alkoxy or C_2-8 alkanoyloxy, A is a C_2-4 alkylene group with a chain length between NH and NHR or $N(O)R'R''$ of at least 2 carbon atoms and R' , R' and R'' are each separately selected from C_1-4 alkyl groups and C_2-4 hydroxyalkyl and C_2-4 dihydroxyalkyl groups in which the carbon atom attached to the nitrogen atom does not carry a hydroxy group and no carbon atom is substituted by two hydroxy groups, or R' and R'' together are a C_2-6 alkylene group which with the nitrogen atom to which R' and R'' are attached forms a heterocyclic group having 3 to 7 atoms in the ring, but with the proviso that at least one of R_1 to R_4 is a group $NH-A-N(O)R'R''$, the compound optionally being in the form of a physiologically acceptable salt.

The compounds of formula (I) contain at least one substituent group $NH-A-N(O)R'R''$ having the terminal, tertiary nitrogen atom in N-oxide form. Although groups of this type are not unknown, for example being described in European Patent Application A-0 145 226 as one of the various alternative forms of substituent in a group of substituted nitroacridones, it had not previously been appreciated that such a substituent confers valuable properties as compared with the compound containing the corresponding group $NH-A-NR'R''$ in which the terminal, tertiary nitrogen atom is not in N-oxide form. Thus, such N-oxides are biochemically activated within neoplastic tissue to form the cytotoxic compound containing

- 3 -

an NH-A-NR'R" group thereby providing the desired anti-cancer activity of this compound but with mitigation of its undesired side effects.

05 It will be seen that in addition to the one or more substituents NH-A-N(O)R'R" in N-oxide form the compounds (I) may contain one or more substituents NH-A-NHR. Whilst these compounds, as compared with those containing no group NH-A-NHR, may exhibit some degree of cytotoxicity and are thus less preferred, this will nonetheless be at a lower level than the corresponding compound in which none of
10 the aminoalkylamino groups is in N-oxide form and full cytotoxicity will only be expressed on conversion of the group(s) NH-A-N(O)R'R" to group(s) NH-A-NR'R".

As regards the groups NH-A-NHR and NH-A-N(O)R'R", A may be branched but is conveniently a straight chain alkylene group,
15 i.e. tetramethylene, especially trimethylene, or particularly ethylene.

R, R' and R" may also have a branched carbon chain but are conveniently straight chain whether they are alkyl groups or hydroxy-substituted alkyl groups. When R, R' or R" is a
20 monohydroxyalkyl group this is conveniently substituted terminally and when R, R' or R" is a dihydroxyalkyl group this is conveniently substituted terminally by one of the hydroxy groups. When R, R' and R" are alkyl the preference is for a group of three or especially two or one carbon atoms and when R, R' and R" are
25 hydroxy-substituted alkyl the preference is for the alkyl group to be of three carbon atoms or, in the case of a monohydroxyalkyl group, alternatively of two carbon atoms. Examples of preferred individual groups R, R' and R" are CH₃, CH₂CH₂CH₃, CH₂CH₂OH, CH₂CH₂CH₂OH, CH(CH₃)CH₂OH, and CH₂CHOHCH₂OH and especially CH₂CH₃.
30 Whilst R' and R" will more usually be identical there can be certain advantages as described hereinafter in having non-identical groups R' and R".


- 4 -

Alternatively, as indicated, R' and R" together with the nitrogen atom to which they are attached may represent a heterocyclic group $-N(CH_2)_n$ where n is 2 to 6, i.e. aziridin-1-yl, azetidin-1-yl, pyrrolidin-1-yl, piperidin-1-yl and perhydroazepin-1-yl, the smaller groups such as azetidin-1-yl and especially aziridin-1-yl being of most interest.

Specific groups NH-A-NHR of particular interest are $NH-(CH_2)_2-NHCH_3$, $NH-(CH_2)_2-NHCH_2CH_2CH_2OH$, $NH-(CH_2)_2-NHCH(CH_3)CH_2OH$, $NH-(CH_2)_2-NHCH_2CHOHCH_2OH$, especially $NH-(CH_2)_2-NHCH_2CH_2OH$ and particularly $NH-(CH_2)_2-NHC_2H_5$, whilst specific groups NH-A-N(O)R'R" of particular interest are $NH-(CH_2)_2-N(O)(CH_3)C_2H_5$, $NH-(CH_2)_2-N(O)(CH_2CH_2OH)_2$, $NH-(CH_2)_2-N(O)(CH_2CH_2CH_2OH)_2$, $NH-(CH_2)_2-N(O)(CH(CH_3)CH_2OH)_2$, $NH-(CH_2)_2-N(O)(CH_2CHOHCH_2OH)_2$, especially $NH-(CH_2)_2-N(O)(CH_3)_2$ and particularly $NH-(CH_2)_2-N(O)(C_2H_5)_2$.

As regards the groups X, the halogeno groups are preferably bromo and especially chloro. Alkoxy and alkanoyloxy groups X may be branched or especially straight chain and are conveniently of 1 or 2 carbon atoms for the alkyl groups and of 2 or 3 carbon atoms for the alkanoyl groups. Examples of such groups X are therefore chloro, amino and especially methoxy, ethoxy, acetyl and propionyl. However hydroxy groups are preferred as the group or groups X.

Formula (II) illustrates the system used for numbering the various positions of the anthracene-9,10-dione ring system.



(II)

- 5 -

It will be seen from this formula that, due to the symmetrical nature of the molecule, certain substitution patterns are equivalent, for example 1,4 and 5,8. Preferences as to positions of substitution are expressed herein by identifying groups R_1 to R_4 which are other than hydrogen in the order R_1 , R_2 , R_3 , R_4 and, as is conventional practice, by identifying substituted positions in the order 1, 2, 3, 4, 5, 6, 7, 8. Thus, for example, the compound having one substituent only at a ring position meta to an oxo group is identified as having that substituent as a group R_1 at position 1 and groups R_2 , R_3 and R_4 which are hydrogen.

Any of positions 1, 2, 3, 4, 5, 6, 7 and 8 in the compound (I) may be substituted by a group $NH-A-N(O)R'R''$ but positions 1, 4, 5 and 8 are of most interest for substitution by such a group and, indeed, also by the other substituent groups. At least one group $NH-A-N(O)R'R''$ in the compound (I) is conveniently at a position meta to an oxo group (so that R_1 is a group $NH-A-N(O)R'R''$ at the 1-position). However, the compound (I) may contain more than one group $NH-A-N(O)R'R''$, which may be different as regards A and/or R' and R'' but which are conveniently identical. Although four or more, particularly three, of such groups may be present, preferred compounds contain two such groups, conveniently with a 1,8, especially a 1,4 and particularly a 1,5 substitution pattern for these groups (so that R_1 is a group $NH-A-N(O)R'R''$ at the 1-position together with either R_2 being a group $NH-A-N(O)R'R''$ at the 4-position or R_3 being a group $NH-A-N(O)R'R''$ at the 5- or 8-position).

The compounds (I) may contain one to three groups $NH-A-NHR$ but conveniently no more than two and preferably no more than one of such groups is present. Where one or more groups $NH-A-NHR$ are present these may differ or not as regards A and/or R between each other and as compared with A and R' and R'' in the group(s) $NH-A-N(O)R'R''$ which are present. Preferably, however, each group $NH-A-NHR$ is identical where more than one of these is present. Where only one group $NH-A-N(O)R'R''$ is present or more than one of such groups which are the same are present, a possibility is for

- 6 -

any group NH-A-NHR to be the same as the group(s) NH-A-N(O)R'R" as regards A and conveniently for R to be the same as R' and/or R". Where one or more groups NH-A-NHR is present there is preferably one such group at a position meta to an oxo group. Compounds
05 containing such a group are conveniently substituted at one or more of positions 1, 4, 5 and 8 by this group or groups. Preferred compounds of this type contain only one group NH-A-NHR and one group NH-A-N(O)R'R", compounds of particular interest having a group NH-A-N(O)R'R" at position 1 and a group NH-A-NHR at position 8
10 or especially at position 5 and particularly at position 4, optionally with substitution by a group or groups X, particularly hydroxy, at one or both of the other of these positions.

Conveniently the compounds (I) contain also at least one and preferably two groups X, particularly hydroxy groups. Once again,
15 a group X may be at any of positions 1, 2, 3, 4, 5, 6, 7 and 8 but such a group may conveniently be at one or two of the positions 1, 4, 5 and 8, providing these are not occupied by a group NH-A-NHR or NH-A-N(O)R'R". Conveniently three groups X may be present when only one group NH-A-N(O)R'R" and no group NH-A-NHR is present and
20 two groups X may be present when two groups NH-A-N(O)R'R" or one group NH-A-N(O)R'R" and one group NH-A-NHR are present, such groups X being the same or different. Preferably position 1 is occupied by a group NH-A-N(O)R'R" and the positions 4, 5 and 8 are occupied by a group X, particularly a hydroxy group in each case, or one of
25 these three positions is occupied by another group NH-A-N(O)R'R" or a group NH-A-NHR and the remaining two positions are occupied by a group X, particularly a hydroxy group in each case. Compounds having a group X at each of positions 5 and 8, particularly a hydroxy group in each case, are preferred, for example those
30 indicated under (1), (5) and (7) below.

Compounds of particular interest thus have one of:

- (1) $R_1 = \text{NH-A-N(O)R'R"}$ (position 1), $R_2 = \text{H}$, $R_3 = R_4 = \text{OH}$ (positions 5 and 8);
(2) $R_1 = \text{NH-A-N(O)R'R"}$ (position 1), $R_2 = \text{OH}$ (position 4),
35 $R_3 = \text{OH}$ (position 5 or position 8), $R_4 = \text{H}$;

- 7 -

- (3) $R_1 = \text{NH-A-N(O)R'R''}$ (position 1), $R_2 = R_3 = R_4 = \text{OH}$ (positions 4, 5 and 8);
- (4) $R_1 = R_3 = \text{NH-A-N(O)R'R''}$ (positions 1 and 8, conveniently being identical groups), $R_2 = R_4 = \text{OH}$ (positions 4 and 5 respectively);
- 05 (5) $R_1 = R_2 = \text{NH-A-N(O)R'R''}$ (positions 1 and 4, conveniently being identical groups), $R_3 = R_4 = \text{OH}$ (positions 5 and 8);
- (6) $R_1 = R_3 = \text{NH-A-N(O)R'R''}$ (positions 1 and 5, conveniently being identical groups), $R_2 = R_4 = \text{OH}$ (positions 4 and 8);
- (7) $R_1 = \text{NH-A-N(O)R'R''}$ at position 1, $R_2 = \text{NH-A-NHR}$ at position 4,
- 10 $R_3 = R_4 = \text{OH}$ at positions 5 and 8;
- (8) $R_1 = \text{NH-A-N(O)R'R''}$ at position 1, $R_2 = \text{OH}$ at position 4, $R_3 = \text{NH-A-NHR}$ at position 5 and $R_4 = \text{OH}$ at position 8, or
- (9) $R_1 = \text{NH-A-N(O)R'R''}$ at position 1, $R_2 = R_3 = \text{OH}$ at positions 4 and 5, and $R_4 = \text{NH-A-NHR}$ at position 8.
- 15 Of these, the compounds of types (1) to (6), particularly of type (5) and especially of type (6) are preferred.

Specific compounds (I) according to the present invention include those compounds of types (1) to (9) just listed in which the or each group NH-A-N(O)R'R'' and any group NH-A-NHR has

20 $\text{A} = (\text{CH}_2)_2$ and R , R' and R'' each separately = CH_3 , $\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{OH}$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$, $\text{CH}(\text{CH}_3)\text{CH}_2\text{OH}$ or $\text{CH}_2\text{CHOHCH}_2\text{OH}$ or particularly CH_2CH_3 . Preferably R' and R'' are identical for each group NH-A-N(O)R'R'' and conveniently where two groups NH-A-N(O)R'R'' are present these are identical. Particularly preferred specific

25 compounds are those of formula (III) and particularly of formula (IV) and their analogues in which the two methyl groups in $\text{N(O)(CH}_3)_2$ are replaced by two n-propyl, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 2,3-dihydroxypropyl or particularly ethyl groups, such compounds being in the free base or



30 a salt form.

A further group of specific compounds, which are also of some interest, consists of compounds analogous to compounds (III) and (IV) and the analogues thereof just mentioned but in which the $\text{NH-(CH}_2)_2\text{-N(O)R'R''}$ group at position 4 or 5 is replaced by a group

35 $\text{NH-(CH}_2)_2\text{-NHCH}_2\text{CH}_2\text{OH}$ or a variant of such a group in which the

- 8 -

2-hydroxyethyl group thereof is replaced by a methyl, n-propyl, 2-hydroxypropyl, 3-hydroxypropyl, 2,3-dihydroxypropyl or particularly an ethyl group.



Certain substituents in the compounds (I) may contain one or
 05 more asymmetric carbon atoms and the compounds will then exist in
 stereoisomeric forms. Moreover, in the case where R' and R" are
 different this will introduce a centre of asymmetry at the nitrogen
 atom in N-oxide form. It will be appreciated that one
 stereoisomeric form of a compound may be of particular interest by
 10 virtue of advantageous physical properties, for example greater
 solubility, or biological activity.

As indicated the compounds (I) may be used in the form of a
 physiologically acceptable salt which will be an acid addition salt
 with an organic or inorganic acid, for example with one of the
 15 acids sulphuric, phosphoric, hydrochloric, hydrobromic, sulphamic,
 citric, lactic, malic, succinic, tartaric, acetic, benzoic,
 gluconic and ascorbic. Although the salts will usually have
 similar physiological properties to the free base they may have the
 advantage of enhanced solubility, etc.

20 The compounds (I) may conveniently be prepared through the
 oxidation of the tertiary amino group(s) of the corresponding
 compound in which each group NH-A-N(O)R'R" in the compound (I) is
 in the form NH-A-NR'R". Thus, for example, anthracene-9,10-diones

- 9 -

containing various [2-(dialkylamino)ethyl]amino, {2-[di-(hydroxy-alkyl)amino]ethyl}amino and [2-(cyclic alkyleneamino)ethyl]amino groups may be oxidized to the ω -N-oxides. Where appropriate the precursor compound which is oxidized may contain one or more

05 modified groups X, R, R' and R" as compared with the parent compound, the groups X, R, R' and R" corresponding to those in the compound (I) being generated after the oxidation has been effected. In particular, it may be appropriate to protect the

10 hydroxy group(s) in groups R, R' and R" which are hydroxyalkyl or dihydroxyalkyl groups or groups X which are hydroxy during the oxidation, for example as an ether group such as benzyloxy, and subsequently regenerate the hydroxy group(s), for example by

15 catalytic reduction of a benzyloxy group. Any suitable oxidizing agent for converting a tertiary aliphatic amine to N-oxide form is suitable, for example hydrogen peroxide, oxone (potassium monopersulphate) and particularly a peracid such as

20 3-chloroperbenzoic acid. Reaction at room temperature in the dark overnight with an excess of such an acid is usually sufficient to effect conversion to the N-oxide.


25 Where the compound (I) can exist in d and l forms as well as the dl form an optically active isomer may be synthesised either substantially free from these other forms, or at least in a major proportion by weight as compared with them, either by using optically active reagents in the synthesis of the compound or,

30 particularly in the case of the optically active compounds in which R' and R" are different, by resolving the dl form, especially by using an optically active inorganic or organic acid to provide two stereoisomeric salts with different physical properties. In such an instance and also where the compound (I) is used in the form of

35 a salt the salt may be prepared by reaction of the organic base (I) with the appropriate inorganic or organic acid according to conventional procedures, usually by simple admixture in solution. The acid addition salts are generally crystalline solids which are relatively soluble in water, methanol, ethanol and similar solvents.

- 10 -

Accordingly the present invention comprises a process for the preparation of a compound of formula (I) as defined hereinbefore which comprises oxidizing a compound of formula (Ia)



in which R^1 , R^2 , R^3 and R^4 correspond to R_1 , R_2 , R_3 and R_4 , respectively, in the compound (I) but with each of the groups of the type $NH-A-N(O)R'R''$ in the compound (I) being instead a group $NH-A-NR'R''$ in the compound (Ia) and one or more groups X , R , R' and R'' in the compound (Ia) optionally instead being in a form convertible to said group or groups present in the compound (I), and where appropriate converting the one or more modified groups X , R , R' and R'' in the compound (Ia) to the form present in the compound (I) and/or forming an acid addition salt with a physiologically acceptable organic or inorganic acid.

Various routes are available for the synthesis of the intermediates which are oxidized to the compounds (I) of the present invention. One very convenient procedure for the preparation of compounds having a group $NH-A-NR'R''$ at the 1 and 4 positions uses the appropriately substituted 2,3-dihydro(leuco)-1,4-dihydroxyanthracene-9,10-dione which is condensed with the appropriate amine $R'R''N-A-NH_2$, the 1,4 positions being activated in the leuco compound for reaction with the amine. Such a condensation may conveniently be effected at a temperature in a range of about 25 or 35 to 50 or 60°C for one or more hours using a solvent such as methanol, ethanol, water, dimethylformamide, 2-methoxyethanol, acetonitrile, nitrobenzene, N,N,N',N' -tetramethylenediamine or mixtures thereof. In some instances a higher temperature and shorter reaction time may be appropriate, for

- 11 -

example with the compounds containing cyclic groups $\text{NR}'\text{R}''$. The leuco derivative is then oxidized to the fully aromatic anthracene-9,10-dione, conveniently using air oxidation or oxidation with hydrogen peroxide, chloranil, sodium perborate or manganese dioxide.

05

Although leuco compounds are primarily of interest for the preparation of intermediates substituted by two $\text{NH-A-NHR}'\text{R}''$ groups, it is possible to use them to prepare compounds containing more than two such groups. Thus by using 2,3-dihydro(leuco)-1,4,5,8-tetrahydroanthracene-9,10-dione and a large excess of an amine $\text{NH-A-NHR}'\text{R}''$ an 8-hydroxyanthracene-9,10-dione having three groups $\text{NH-A-NHR}'\text{R}''$ at the 1, 4 and 5 positions may be prepared.

10

The leuco derivatives themselves are obtainable by heat treatment of the corresponding fully aromatic 1,4-dihydroxyanthracene-9,10-dione, conveniently by heating at above 90°C for 1 hour or more in a stream of nitrogen and, if necessary, in the presence of a suitable reducing agent such as sodium dithionite or zinc dust. Various anthracene-9,10-diones, particularly hydroxyanthracene-9,10-diones, are commercially available and various syntheses for such compounds are also reported in the literature. One suitable procedure for their preparation involves the reaction of an appropriately substituted phthalic anhydride with hydroquinone in the presence of aluminium chloride and sodium hydroxide at 180°C for one hour or more. Anthracene-9,10-diones containing one form of substituent X can be modified to provide other forms of substituent X so that, for example, a dione containing an amino group can be treated with sodium hydroxide/dithionite to yield the corresponding hydroxy substituted compound.

15

20

25

30

35

Other suitable procedures for the preparation of intermediates for oxidation to the N-oxide include the reaction of the appropriate chloro substituted anthracene-9,10-dione with the appropriate amine $\text{R}'\text{R}''\text{N-A-NH}_2$, for example by heating with a excess of the amine at its reflux temperature for one or more hours. Certain of these chloroanthracene-9,10-diones are known and

- 12 -

various syntheses for such compounds are also reported in the literature. Thus, for example, 1,5-dichloro-4,8-dihydroxy-anthracene-9,10-dione may be prepared by selective chlorination of 1,4,5,8-tetrahydroxyanthracene-9,10-dione using a stoichiometric amount of sulphuryl chloride and controlled temperature. This precursor may then be used to prepare an intermediate having groups NH-A-NR'R" at the 1 and 5 positions and hydroxy groups at the 4 and 8 positions, the hydroxy groups conveniently being protected during the reaction with the amine R'R"N-A-NH₂. A similar approach is suitable for the preparation of other chlorohydroxyanthracene-9,10-dione intermediates.

Where the compound (I) contains one or more groups NH-A-NHR in addition to the one or more groups NH-A-NR'R" the compound may conveniently be produced by reacting a suitable precursor as discussed above with a mixture of amines RN-A-NH₂ and R'R"N-A-NH₂, the resultant mixture of products then being separated, for example by chromatography. Thus, for example, 2,3-dihydro(leuco)-1,4-dihydroxyanthracene-9,10-dione on reaction with a mixture of 2-(2-hydroxyethylamino)ethylamine and 2-(diethylamino)ethylamine will yield a mixture of 1,4-bis{[2-(diethylamino)ethyl]amino}-anthracene-9,10-dione, 1,4-bis{[2-(2-hydroxyethylamino)ethyl]amino}-anthracene-9,10-dione and 1-{2-(diethylamino)ethyl}amino-4-{[2-(2-hydroxyethylamino)ethyl]amino}anthracene-9,10-dione from which the last mentioned compound may be separated, for example by chromatography. On oxidation, only the tertiary nitrogen atom of the [2-(diethylamino)ethyl]amino group will be converted to N-oxide form.

Where one or more substituents X is present it may be appropriate, depending on the route of synthesis, to have these present throughout in their final form or to generate the desired groups at a later stage in the synthesis. Ether and ester groups X may of course readily be prepared by modification of hydroxy groups according to known procedures, precursors containing a hydroxy group X more often being described in the literature than those containing a corresponding ether or ester substituent.

- 13 -

It will be appreciated, however, that various alternative methods for the preparation of the compounds (I) and intermediates therefor may be used as will be apparent in particular from the literature relating to such intermediates. Further details of the preparation of intermediates for the preparation of the compounds (I) of the present invention are to be found in U.S. Patent 4,197,249 and U.K. Patent GB 2,004,293B referred to hereinbefore.

Certain of the intermediates corresponding to compounds (I) described herein but without the tertiary amine group(s) in N-oxide form are novel and are within the scope of this invention. Such intermediates include particularly those of formula (Ia) in which at least one of R_1 to R_4 is a group $NH-A-NR'R''$ and at least one other is a different group $NH-A-NR'R''$ or a group $NH-A-NHR$.

The compounds (I) may be formulated with a physiologically acceptable diluent or carrier for use as pharmaceuticals for both veterinary, for example in mammals, and particularly human use by a variety of methods. For instance, they may be applied as a composition incorporating a liquid diluent or carrier, for example an aqueous or oily solution, suspension or emulsion, which may often be employed in injectable form for parenteral administration and therefore may conveniently be sterile and pyrogen free. Oral administration may also be used and although compositions for this purpose may incorporate a liquid diluent or carrier, it is more usual to use a solid, for example a conventional solid carrier material such as starch, lactose, dextrin or magnesium stearate. Such solid compositions may take the form of powders but are more conveniently of a formed type, for example as tablets, cachets, or capsules (including spansules). Alternative, more specialized types of formulation include liposomes and nanoparticles.

Other types of administration than by injection or through the oral route which are of use in both human and veterinary contexts include the use of suppositories or pessaries. Another form of pharmaceutical composition is one for buccal or nasal administration or alternatively drops for administration into the eye which may

- 14 -

conveniently contain a sterile liquid diluent or carrier. Other formulations for topical administration include lotions, ointments, creams, gels and sprays.

05 Compositions may be formulated in unit dosage form, i.e. in the form of discrete portions containing a unit dose, or a multiple or sub-unit of a unit dose.

10 Whilst the dosage of the compound used will vary according to the activity of the particular compound and the condition being treated, it may be stated by way of guidance that a dosage selected in the range from 0.1 to 20 mg/kg per body weight per day, particularly in the range from 0.1 to 5 mg/kg of body weight per day, will often be suitable although higher doses than this, for example in the range from 0.1 to 50 mg/kg of body weight per day (or possibly even as high as described in U.S. Patent 4,197,249) 15 may be considered in view of the lower level of toxic side effects obtained with the compounds (I). This dosage regime may be continued for however many days is appropriate to the patient in question, the daily dosages being divided into several separate administrations if desired. Thus, for example, in the case of 20 conditions such as advanced breast cancer, non-Hodgkin's lymphoma and hepatoma, treatment for one day followed by a repeated dose after an interval, such as 21 days, may be appropriate whilst for the treatment of acute non-lymphocytic leukaemia, treatment over 5 consecutive days may be more suitable.

25 The compounds (I) are of particular value for the treatment of cancer in warm blooded animals including humans. The compounds are of interest in relation to the treatment of solid tumours, such as various forms of sarcoma and carcinoma, and also for disseminated tumours such as leukaemias. Areas of particular interest are the 30 treatment of non-Hodgkins lymphoma, of breast cancer, and of acute non-lymphocytic leukaemia. In the treatment of cancer parenteral and sometimes topical administration is often of particular interest. Moreover, it may be advantageous to use the compounds (I) in a combined treatment, given separately or together in the same 35 composition, with other anti-cancer agents, such as mitotic

- 15 -

inhibitors, for example vinblastine; alkylating agents, for example cis-platin, carboplatin and cyclophosphamide; other antimetabolites, for example 5-fluorouracil, cytosine arabinoside and hydroxyurea; intercalating antibiotics, for example adriamycin and bleomycin; enzymes, for example asparaginase; topoisomerase inhibitors, for example etoposide and biological response modifiers, for example interferon.

In a variation of the usual procedure which relies upon the anaerobic conditions within neoplastic tissue to effect selective reduction of the N-oxide in such tissue, the selectivity as between neoplastic and normal tissue can be enhanced. Thus antibodies can be raised against tumours by conventional procedures, particularly using hybridoma technology, and linked covalently to a reductase using one of various conventional linking agents. The conjugate is administered to the patient when it localises in the body at the tumour site and the compound (I) is then administered, the action of the reductase enhancing the specificity of the action of the compound at the tumour site.

The present invention thus includes a method suitable for aiding regression and palliation of cancer which comprises administering to a patient a therapeutically effective amount of a compound (I) as defined hereinbefore.

In addition to their anti-cancer use the compounds (I) are of interest for various other pharmaceutical applications in view of their activity as chelating agents.

The invention is illustrated by the following Examples.

EXAMPLES

Example 1

Preparation of 1,5-bis-[[2-(diethylamino-N-oxide)ethyl]amino]anthracene-9,10-dione

(1) 1,5-bis-[[2-(diethylamino)ethyl]amino]anthracene-9,10-dione.

A solution of 5.52g (0.02 mol) of 1,5-dichloroanthracene-9,10-dione in 23.2g (0.2 mol) of 2-(diethylamino)ethylamine is heated at reflux temperature for 4 hours. The mixture is cooled in an

- 16 -

ice-bath and 100 ml concentrated hydrochloric acid is added with stirring. This acidic mixture is extracted with three aliquots of 200 ml of diethylether followed by three aliquots of 200 ml of chloroform. The aqueous layer is collected and made alkaline with sodium hydroxide solution, extracted into chloroform and evaporated in vacuo at 30°C. The oily residue is washed with water, dissolved in methanol and titrated with hydrogen chloride in dry diethyl ether to give a precipitate which is dried to yield 5.2 g of the title compound as the hydrochloride in the form of a dark red solid, m.p. 158.5-159.5°C; λ_{\max} (deionised water) (E/cm/M) 231 nm (34280), 516 nm (10690).

(2) 1,5-bis-[[2-(diethylamino-N-oxide)ethyl]amino]anthracene-9,10-dione.

A solution of 1.25 g (0.004 mol) of 1,5-[[2-(diethylamino)-ethyl]amino]anthracene-9,10-dione free base (prepared by washing the oily residue of (1) with water and drying) in 20 ml of chloroform is cooled in an ice bath while stirring. This solution is treated with 1.2 g of 3-chloroperbenzoic acid, allowed to come to room temperature and left for 18 hours protected from light. The mixture is evaporated in vacuo to a small volume and is subject to flash chromatography using a column of 70-230 mesh (60 A) silica gel and an eluting solvent of chloroform:methanol (5:1 v/v) to yield 0.45g of the title compound, m.p. (following recrystallisation from chloroform to give dark red crystals) 125-129°C (decomposition); λ_{\max} (deionised water) (E/cm/M) 231 nm (44030), 516 nm (11330).

Example 2

Preparation of 1,4-bis-[[2-(diethylamino-N-oxide)ethyl]amino]-anthracene-9,10-dione.

(1) 1,4-bis-[[2-(diethylamino)ethyl]amino]anthracene-9,10-dione.

A mixture of 5 g (0.021 mol) of 1,4-dihydroxyanthracene-9,10-dione and 2 g (0.014 mol) of sodium dithionite in 20 ml water is stirred whilst heating under nitrogen at 90°C until the mixture

- 17 -

turns from orange to brown indicating the presence of 2,3-dihydro-(leuco)-1,4-dihydroxyanthracene-9,10-dione. To this reaction mixture is added dropwise 20 g (0.17 mol) of 2-(diethylamino)-ethylamine over a 30 minute period. The mixture is heated at 50-55°C for 2 hours and 20 ml of ethanol are added. The solution is then aerated and 200 ml of 2M hydrochloric acid are added. The acidic mixture is washed with 3 x 200 ml of diethylether followed by 3 x 200 ml of chloroform. The aqueous phase is made alkaline by the addition of sodium hydroxide solution and extracted with 3 x 200 ml chloroform. The mixture is evaporated in vacuo to a 5 ml volume and subjected to flash chromatography using a column of 70-230 mesh (60 A) silica gel (SiO₂) and an eluting solvent of chloroform followed by chloroform:methanol (1:1 v/v). The blue black material removed from the column in chloroform/methanol is evaporated in vacuo to yield 0.7 g of the title compound, m.p. 107.5-108.5°C; λ_{\max} (deionised water) (E/cm/M) 256 nm (50950), 584 nm (18650). (2) 1,4-bis-[[2-(diethylamino-N-oxide)ethyl]amino]anthracene-9,10-dione.

To an ice-cool solution of 0.8 g (0.0018 mol) of 1,4-[[2-(diethylamino)ethyl]-amino]anthracene-9,10-dione in 15 ml of dry chloroform is added 1.5 g (0.008 mol) of 3-chloroperbenzoic acid. The mixture is stirred for 30 minutes, allowed to come to room temperature and then left for 18 hours protected from light. The mixture is evaporated in vacuo to a 5 ml volume and subjected to reverse phase chromatography using a column of octadecylsilane bonded to silica gel, 10 μ M particle size, and methanol:ammonium formate, 0.5M, pH 4.25 (30:70 v/v), as an eluting solvent. The eluate fractions identified by thin-layer chromatography (chloroform/methanol 1:1 v/v) to contain a single dark blue-coloured component are pooled, extracted with chloroform and evaporated in vacuo. Drying over phosphorous pentoxide yields 0.17 g of the title compound as a dark blue solid, m.p. 115-118°C (decomposition); λ_{\max} (deionised water) (E/cm/M) 256 nm (20900), 584 nm (5880).

- 18 -

Example 3Preparation of 1-[[2-(diethylamino-N-oxide)ethyl]amino]anthracene-9,10-dione.(1) 1-[[2-(diethylamino)ethyl]amino]anthracene-9,10-dione.

05 A mixture of 12.1 g (0.05 mol) of 1-chloroanthraquinone and 58 g (0.5 mol) of 2-(diethylamino)ethylamine is heated under reflux for 2 hours. The reaction product is worked up as described in Example 1 except that the oily residue is only washed with water and then dried to yield 3.24 g of the title compound as the free
10 base in the form of an orange-red solid, m.p. 98°C; λ_{\max} (deionised water) (E/cm/M) 248 nm (27180), 493 nm (5770).

(2) 1-[[2-(diethylamino-N-oxide)ethyl]amino]anthracene-9,10-dione.

1.25 g (0.0039 mol) of 1-[[2-(diethylamino)ethyl]amino]anthracene-9,10-dione and 1.2 g (0.0068M) 3-chloroperbenzoic acid
15 are reacted for 18 hours as described in Example 1. Silica gel flash chromatography as in Example 1 but with chloroform/methanol (1:5 v/v) as eluting solvent gives an eluate which is evaporated in vacuo. Recrystallisation of the residue from chloroform yields
20 0.45 g of the title compound as an orange-red solid, m.p. 120-125°C (decomposition); λ_{\max} (deionised water) (E/cm/M) 248 nm (32330), 493 nm (6180).

Example 4Preparation of 1,8-bis-[[2-(diethylamino-N-oxide)ethyl]amino]anthracene-9,10-dione.(1) 1,8-bis-[[2-(diethylamino)ethyl]amino]anthracene-9,10-dione.

25 A mixture of 5.52 g (0.02 mol) of 1,8-dichloroanthraquinone and 23.2 g (0.2M) 2-(diethylamino)ethylamine is heated under reflux for 3 hours. The reaction product is washed up as described in Example 1 except that the oily residue is only washed with water
30 and then dried to yield 3.63 g of the title compound as the free base in the form of a purple solid, m.p. 103.5-106.5°C;
 λ_{\max} (deionised water) (E/cm/M) 236 nm (54940), 542 nm (13200).

- 19 -

(2) 1,8-bis-[[2-(diethylamino-N-oxide)ethyl]amino]anthracene-9,10-dione.

1.25 g (0.003 mol) of 1,8-[[2-(diethylamino)ethyl]amino]-anthracene-9,10-dione and 2.5 g (0.0135 mol) 3-chloroperbenzoic acid are reacted for 4 hours as described in Example 1. Flash chromatography as in Example 1 but using chloroform/methanol (1:1 v/v) eluting solvent gives an eluate which is evaporated in vacuo. Recrystallisation of the residue from chloroform yields 0.18 g of the title compound as a purple solid
m.p. 118-119°C (decomposition); λ_{\max} (deionised water) (E/cm/M) 236 nm (39530), 542 nm (8360).

Example 5

Preparation of 1,4-bis-[[2-(diethylamino-N-oxide)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

(1) 1,4-bis-[[2-(diethylamino)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

A mixture of 1.0 g (0.0037 mol) of leuco-1,4,5,8-tetrahydroxy-anthraquinone and 5 g (0.043 mol) of N,N-diethylaminoethylamine in methanol is heated for 10 minutes to reflux temperature. The ethanol and unreacted N,N-diethylaminoethylamine are removed by distillation in vacuo and the remaining solid recrystallised from ethyl acetate to yield 0.64 g of the title compound as a dark blue solid, m.p. 203°C as the dihydrochloride salt; λ_{\max} (distilled water) (E/cm/M) 239 nm (44910), 605 nm (10890), 659 nm (10710).

(2) 1,4-bis-[[2-(diethylamino-N-oxide)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

0.108 g (0.00023 mol) of 1,4-bis-[[2-(diethylamino)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione is dissolved in 5 ml dichloromethane and the solution is cooled in an ice-bath whilst stirring. The solution is treated with 0.2 g (0.000115 mol) of 3-chloroperbenzoic acid, allowed to come to room temperature and left for 18 hours protected from light. The mixture is then subjected to flash column chromatography using a column of silica gel (60A) and a step-gradient eluting solvent of

- 20 -

dichloromethane:methanol:triethylamine starting with dichloromethane:triethylamine (99:1 v/v) followed by dichloromethane:methanol:triethylamine (90:9:1 v/v/v) then
05 finally methanol:triethylamine (99:1 v/v). The major eluting blue fraction is collected, filtered and evaporated in vacuo to yield 0.08 g of the title compound as a dark blue solid, m.p. 155-158°C (decomposition); λ_{\max} (distilled water) (E/cm/M) 240 nm (15690), 609 nm (8809), 662 nm (5750).

10 Example 6Preparation of 1,4-bis-[[2-(dimethylamino-N-oxide)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione(1) 1,4-bis-[[2-(dimethylamino)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

15 A solution of 1.0 g (0.0037 mol) of leuco-1,4,5,8-tetrahydroxyanthraquinone in methanol is refluxed under nitrogen. To this is slowly added 2 g (0.023 mol) of N,N-dimethylaminoethylamine and the mixture is stirred at 50°C for 1 hour. The mixture is then stirred for 16 hours in air. The ethanol and unreacted N,N-dimethylaminoethylamine are removed by distillation in vacuo and the remaining
20 solid is recrystallised from ethyl acetate/methanol (2:1 v/v) to yield 0.68 g of the title compound as a dark blue solid, m.p. 199-200.5°C as the dihydrochloride salt, λ_{\max} (distilled water) (E/cm/M) 242 nm (43270), 606 nm (18050), 658 nm (16220).

25 (2) 1,4-bis-[[2-(dimethylamino-N-oxide)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

A solution of 0.10 g (0.000242 mol) of 1,4-bis-[[2-(dimethylamino)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione in 5 ml dichloromethane is cooled in an ice-bath whilst stirring. To this
30 solution is added 0.2 g (0.000115 mol) of 3-chloroperbenzoic acid and the product is allowed to come to room temperature and left for 8 hours protected from light. The mixture is then subjected to flash column chromatography using a column of

- 21 -

silica gel (60A) and a step-gradient eluting solvent of dichloromethane:methanol:triethylamine starting with dichloromethane:methanol (50:50 v/v) followed by dichloromethane:methanol (20:80 v/v) and finally methanol:triethylamine (99:1 v/v). The last eluting blue fraction is collected, filtered and evaporated in vacuo to yield 0.075 g of the title compound as a dark blue solid, m.p. 124-128°C (decomposition); λ_{\max} (distilled water) (E/cm/M) 222 nm (15500), 612 nm (6110), 664 nm (3685).

10 Example 7

Preparation of 1,4-bis-[[2-(diethylamino-N-oxide)propyl]amino]-5,8-dihydroxyanthracene-9,10-dione

(1) 1,4-bis-[[2-(diethylamino)propyl]amino]-5,8-dihydroxyanthracene-9,10-dione

15 A mixture of 1.0 g (0.0037 mol) of leuco-1,4,5,8-tetrahydroxy-anthraquinone in 20 ml of aqueous potassium carbonate (5% w/v) and 0.2 g of sodium dithionite is stirred and flushed with nitrogen. The mixture is then treated with 4 g (0.003 mol) of N,N-diethylaminopropylamine and stirred at 80°C for 18 hours in
20 air. The ethanol and unreacted N,N-diethylaminopropylamine are removed by distillation in vacuo and the remaining solid is recrystallised from ethyl acetate to yield 0.8 g of the title compound as a dark blue solid, m.p. 126-128°C; λ_{\max} (distilled water) (E/cm/M) 241 nm (15370), 611 nm (12460), 668 nm (11280).

25 (2) 1,4-bis-[[2-(diethylamino-N-oxide)propyl]amino]-5,8-dihydroxyanthracene-9,10-dione

A solution of 0.10 g (0.000242 mol) of 1,4-bis-[[2-(diethylamino)propyl]amino]-5,8-dihydroxyanthracene-9,10-dione is dissolved in 5 ml dichloromethane and cooled in an ice-bath whilst stirring.
30 The solution is treated with 0.2 g (0.000115 mol) of 3-chloroperbenzoic acid, allowed to come to room temperature and left for 8 hours protected from light. The mixture is then subjected to flash column chromatography using a column of silica gel (60A) and a step-gradient eluting solvent of

- 22 -

dichloromethane:methanol:triethylamine starting with dichloromethane:methanol (50:50 v/v) followed by dichloromethane:methanol (20:80 v/v) and finally methanol:triethylamine (99:1 v/v). The last eluting blue fraction
05 is collected, filtered and evaporated in vacuo to yield 0.8 g of the title compound as a dark blue solid which remains as a solid in a dessicator but becomes sticky on attempting a melting point determination; λ_{\max} (distilled water) (E/cm/M) 248 nm (14220), 614 nm (8600), 666 nm (5144).

10 Example 8

Preparation of 1-[[2-(diethylamino-N-oxide)ethyl]amino]-4-[[2-[(2-hydroxyethyl)amino]ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione
(1) 1-[[2-(diethylamino)ethyl]amino]-4-[[2-[(2-hydroxyethyl)-amino]ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

15 A solution of 1.0 g (0.0037 mol) of leuco-1,4,5,8-tetrahydroxy-anthraquinone in 10 ml ethanol is heated at 50°C under nitrogen with 2.1 g (0.00184 mol) of N,N-diethylaminoethylamine and 1.9 g (0.00184 mol) of 2-hydroxyethylaminoethylamine for 6 hours and then stirred in air for 16 hours. The ethanol and unreacted
20 N,N-diethylaminoethylamine and 2-hydroxyethylaminoethylamine are removed by distillation in vacuo. The remaining solid is dissolved in dichloromethane:methanol:0.3% w/v aqueous ammonia (96:3:1 v/v/v) and subjected to flash chromatography using silica gel (60A) and dichloromethane:methanol:aqueous ammonia (96:3:1 v/v/v) as eluting
25 solvent. The major eluting fraction is evaporated in vacuo and the solid again column chromatographed on silica gel (60A) using a step gradient of dichloromethane:methanol (50:50 v/v) followed by dichloromethane:methanol:aqueous ammonia (49.75:49.75:0.5 v/v/v). The major eluting blue fraction is collected, filtered and
30 evaporated in vacuo to yield 0.2 g of the title compound as a dark blue solid, m.p. 165-167°C; λ_{\max} (distilled water) (E/cm/M) 232 nm (33850), 608 nm (20782), 660 nm (18900).

- 23 -

(2) 1-[[2-(diethylamino-N-oxide)ethyl]amino]-4-[[2-[(2-hydroxyethyl)amino]ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

A solution of 0.10 g (0.00023 mol) of 1-[[2-(diethylamino)-ethyl]amino]-4-[[2-[(2-hydroxyethyl)amino]ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione is dissolved in 5 ml dichloromethane and cooled in an ice-bath whilst stirring. This solution is treated with 0.1 g (0.000058 mol) of 3-chloroperbenzoic acid, allowed to come to room temperature and left for 18 hours protected from light. The mixture is then subjected to flash column chromatography using a column of silica gel (60A) and a step-gradient eluting solvent as described for Example 6. The last eluting blue fraction is collected, filtered and evaporated in vacuo to yield 0.06 g of the title compound as a dark blue solid, m.p. 92-93.5°C (decomposition); λ_{max} (phosphate buffer pH 7.4) (E/cm/M) 238 nm (8203), 607 nm (6396), 658 nm (4733).

Example 9

Preparation of 1-[[2-(diethylamino-N-oxide)ethyl]amino]-4-[[2-(ethylamino)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

(1) 1-[[2-(diethylamino)ethyl]amino]-4-[[2-(ethylamino)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

A suspension of 1.0 g (0.0037 mol) of leuco-1,4,5,8-tetrahydroxyanthraquinone in 10 ml of propan-2-ol is stirred at 50°C under nitrogen. To the mixture is added 2.1 g (0.00184 mol) of N,N-diethylaminoethylamine and 1.6 g (0.00184 mol) of N-ethylaminoethylamine the product is stirred in for 12 hours under nitrogen and then for a further 6 hours in air. The ethanol and unreacted N,N-diethylaminoethylamine and N-ethylaminoethylamine are removed by distillation in vacuo and the remaining solid is recrystallised from ethyl acetate to yield 0.2 g of the title compound as a dark blue solid, m.p. 161-163°C; λ_{max} (distilled water) (E/cm/M) 242 nm (15520), 609 nm (10196), 659 nm (12640).

- 24 -

(2) 1-[[2-(diethylamino-N-oxide)ethyl]amino]-4-[[2-(ethylamino)-ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

0.10 g (0.00023 mol) of 1-[[2-(diethylamino)ethyl]amino]-4-[[2-(ethylamino)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione is dissolved in 5 ml dichloromethane and the solution is cooled in an ice-bath whilst stirring. To the solution is added 0.1 g (0.000058 mol) of 3-chloroperbenzoic acid and it is then allowed to come to room temperature and left for 16 hours protected from light. The mixture is then subjected to flash column chromatography using a column of silica gel (60A) and a step-gradient eluting solvent of dichloromethane:methanol (90:10 v/v) followed by dichloromethane:methanol:0.3% w/v aqueous ammonia (90:9:1 v/v/v). The last eluting blue fraction is collected, filtered and evaporated in vacuo to yield 0.08 g of the title compound as a dark blue solid, which remains as a solid in a dessicator but becomes sticky on attempting a melting point determination; λ_{\max} (distilled water) (E/cm/M) 244nm (14776), 612 nm (10080), 662 nm (6532).

Example 10

20 Preparation of 1,4-bis-[[2-(di(2-hydroxyethyl)amino)-N-oxide ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

(1) 1,4-bis-[[2-(di(2-hydroxyethyl)amino)ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

A mixture of 1.0 g (0.0037 mol) of leuco-1,4,5,8-tetrahydroxy-anthraquinone and 4 g (0.003 mol) of N,N-di(2-hydroxyethyl)amino-ethylamine treated as described in Example 7 produces a dark solid. This is dissolved in 10 ml of methanol:dichloromethane (50:50 v/v) subjected to silica gel (60A) flash chromatography using a step gradient of dichloromethane:methanol (90:10 v/v) followed by dichloromethane:methanol (50:50 v/v). The major eluting fraction is collected, filtered and evaporated in vacuo to yield the title compound as a dark blue solid, which remains as a solid in a dessicator but becomes sticky on attempting a melting point determination; λ_{\max} (distilled water) (E/cm/M) 235 nm (12726), 605 nm (7236), 655 nm (6396).

- 25 -

(2) 1,4-bis-[[2-(di(2-hydroxyethyl)amino)ethyl]amino]-5,8-dihydroxy-anthracene-9,10-dione

0.09 g (0.00017 mol) of 1,4-bis-[[2-(di(2-hydroxyethyl)ethyl]-amino]-5,8-dihydroxyanthracene-9,10-dione is dissolved in 5 ml
05 dichloromethane and the solution is cooled in an ice-bath whilst stirring. To the solution is added 0.1 g (0.00044 mol) of 3-chloroperbenzoic acid and it is then allowed to come to room temperature and left for 16 hours protected from light. The reaction mixture is filtered and the solid washed five times
10 with 25 ml aliquots of dichloromethane followed five times by 10 ml aliquots of methanol. The residue is filtered to yield 0.03 g of the title compound as a dark blue solid, m.p. 134.5 - 135.5°C; λ_{\max} (distilled water) (E/cm/M) 238 nm (24527), 610 nm (9331), 666 nm (5435).

15 Example 11

Preparation of 1-[[2-(dimethylamino-N-oxide)ethyl]amino]-4-[[2-[2-hydroxyethyl]amino]ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione.

(1) 1-[[2-(dimethylamino)ethyl]amino]-4-[[2-[2-hydroxyethyl]amino]ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione

A mixture of 1.0 g (0.0037 mol) of leuco-1,4,5,8-tetrahydroxy-anthraquinone, 1.6g (0.00184 mol) of N,N-dimethylaminoethylamine and 1.9g(0.00184 mol) of 2-hydroxyethylaminoethylamine in ethanol is stirred under nitrogen at room temperature for 3 hours. The mixture
25 is then stirred for 16 hours in air. The ethanol and unreacted N,N-dimethylaminoethylamine and 2-hydroxyethylaminoethylamine are removed by distillation in vacuo. The resulting solid is dissolved in dichloromethane:methanol:0.3% w/v aqueous ammonia (49.75: 49.75:0.5 v/v/v) and subjected to column chromatography on
30 silica gel (60A). The chromatography procedure is repeated using dichloromethane:methanol:triethylamine (90:9:1 v/v/v) and the major eluting fraction is collected, filtered and evaporated in vacuo to yield 0.260g of the title compound as a dark blue solid, m.p. 136-140°C as the dihydrochloride; λ_{\max} (distilled water) (E/cm/M) 244nm (30430), 607nm (15683), 658nm (13560).
35

- 26 -

(2) 1-[[2-(dimethylamino-N-oxide)ethyl]amino]-4-[[2-[2-hydroxyethyl)amino]ethyl]amino]-5,8-dihydroxyanthracene-9,10-dione.

0.10g(0.00023 mol) of 1-[[2-(dimethylamino)ethyl]amino]-4-[[2-
05 [(2-hydroxyethyl)amino]ethyl]amino]-5,8-
dihydroxyanthracene-9,10-dione is dissolved in 5ml of
dichloromethane and the solution is cooled in an ice-bath whilst
stirring. To this solution is added 0.1g (0.000058 mol) of
3-chloroperbenzoic acid and it is then allowed to come to room
10 temperature and left for 18 hours protected from light. The
mixture is then treated as described in Example 8 to yield 0.03g of
the title compound as a dark blue solid, m.p. 128-132°C
(decomposition); λ_{\max} (phosphate buffer pH 7.4)
(E/cm/M)240nm(13243), 610nm(6475), 664nm(5915).

15 Example 12 : Biological Activity

The cytotoxicity was compared of the four compounds

- (a) 1,8-bis-[[2-(diethylamino-N-oxide)ethyl]amino]anthracene-9,10-
dione
(b) 1,4-bis-[[2-(diethylamino-N-oxide)ethyl]amino]anthracene-9,10-
20 dione,
(c) 1-[[2-(diethylamino-N-oxide)ethyl]amino]anthracene-9,10-dione,
and
(d) 1,5-bis-[[2-(diethylamino-N-oxide)ethyl]amino]anthracene-9,10-
dione,
25 under aerobic and anaerobic conditions and a comparison was also
made with the cytotoxicity of the corresponding compound in which
the terminal, tertiary nitrogen atom is not in N-oxide form.

Cells of the MCF-7 human breast cancer cell line (5×10^5 /ml)
were seeded into 12-well culture plates containing RPMI 1640 medium
30 (Flow Labs., Irving, Scotland) supplemented with 10% v/v foetal
bovine serum and grown at 37°C to confluence. In the aerobic
experiments, the cells were treated with one of a range of
concentrations of the compound and then incubated in air
for 24 hours. In the anaerobic experiments, following treatment of

- 27 -

the cells with compound the culture plates were placed in a gas-tight chamber (Flow Labs) which was flushed with nitrogen for 30 minutes, the cells then being incubated for 24 hours under nitrogen. Following both types of experiment the cells were washed
05 free of the compound with isotonic saline and were then grown for a further 3 days in air. The surviving monolayer cells were counted using a Coulter counter.


The results obtained are presented in the Figure, the different parts (a) to (d) of which correspond to the four compounds
10 identified above by these letters. In each part of the Figure plot 1 corresponds to the N-oxide under aerobic conditions, plot 2 corresponds to the N-oxide under anaerobic conditions and plot 3 corresponds to the corresponding tertiary amine under anaerobic conditions.

15 The Figure shows that each of the N-oxides is substantially non-cytotoxic in air but these compounds are cytotoxic when cells are exposed to them in nitrogen, although the level of cell kill observed is not necessarily fully equivalent to the level resulting from the use of an equimolar amount of the parent tertiary amines
20 under the same conditions.

- 28 -

CLAIMS

1. A compound of formula (I)



(I)

in which R_1 , R_2 , R_3 and R_4 are each separately selected from hydrogen, X, NH-A-NHR and NH-A-N(O)R'R'' wherein X is hydroxy, halogeno, amino, C_{1-4} alkoxy or C_{2-8} alkanoyloxy, A is a C_{2-4} alkylene group with a chain length between NH and NHR or N(O)R'R'' of at least 2 carbon atoms and R, R' and R'' are each separately selected from C_{1-4} alkyl groups and C_{2-4} hydroxyalkyl and C_{2-4} dihydroxyalkyl groups in which the carbon atom attached to the nitrogen atom does not carry a hydroxy group and no carbon atom is substituted by two hydroxy groups, or R' and R'' together are a C_{2-6} alkylene group which with the nitrogen atom to which R' and R'' are attached forms a heterocyclic group having 3 to 7 atoms in the ring, but with the proviso that at least one of R_1 to R_4 is a group NH-A-N(O)R'R'' , the compound optionally being in the form of a physiologically acceptable salt.

2. A compound according to Claim 1, in which R_1 , R_2 , R_3 and R_4 are each separately selected from hydrogen, hydroxy, NH-A-NHR and NH-A-N(O)R'R'' .
3. A compound according to Claim 1 or 2, in which A is ethylene.
4. A compound according to Claim 1, 2 or 3, in which R, R' and R'' are each separately selected from CH_3 , CH_2CH_3 , $\text{CH}_2\text{CH}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{OH}$, $\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$, $\text{CH}(\text{CH}_3)\text{CH}_2\text{OH}$ and $\text{CH}_2\text{CHOHCH}_2\text{OH}$.


- 29 -

5. A compound according to any of the preceding claims, which contains one or two groups NH-A-N(O)R'R'' , these being separately selected from $\text{NH-(CH}_2)_2\text{-N(O)(CH}_3)_2$, $\text{NH-(CH}_2)_2\text{-N(O)(CH}_3)_2\text{C}_2\text{H}_5$, $\text{NH-(CH}_2)_2\text{-N(O)(C}_2\text{H}_5)_2$, $\text{NH-(CH}_2)_2\text{-N(O)(CH}_2\text{CH}_2\text{OH)}_2$,
 05 $\text{NH-(CH}_2)_2\text{-N(O)(CH}_2\text{CH}_2\text{CH}_2\text{OH)}_2$, $\text{NH-(CH}_2)_2\text{-N(O)CH(CH}_3\text{)OH}$ and $\text{NH-(CH}_2)_2\text{-N(O)(CH}_2\text{CHOHCH}_2\text{OH)}_2$.
6. A compound according to any of the preceding claims, which contains one group NH-A-N(O)R'R'' and one group NH-A-NR , this latter group being selected from $\text{NH-(CH}_2)_2\text{-NHCH}_3$, $\text{NH-(CH}_2)_2\text{-NHC}_2\text{H}_5$,
 10 $\text{NH-(CH}_2)_2\text{-NHCH}_2\text{CH}_2\text{OH}$, $\text{NH-(CH}_2)_2\text{-NHCH}_2\text{CH}_2\text{CH}_2\text{OH}$, $\text{NH-(CH}_2)_2\text{-NHCH(CH}_3\text{)CH}_2\text{OH}$ and $\text{NH-(CH}_2)_2\text{-NHCH}_2\text{CHOHCH}_2\text{OH}$.
7. A compound according to any of Claims 1 to 5, in which
 (1) $\text{R}_1 = \text{NH-A-N(O)R'R''}$ at position 1, $\text{R}_2 = \text{H}$, $\text{R}_3 = \text{R}_4 = \text{OH}$ at positions 5 and 8;
 15 (2) $\text{R}_1 = \text{NH-A-N(O)R'R''}$ at position 1, $\text{R}_2 = \text{OH}$ at position 4, $\text{R}_3 = \text{OH}$ at position 5 or position 8 and $\text{R}_4 = \text{H}$;
 (3) $\text{R}_1 = \text{NH-A-N(O)R'R''}$ at position 1 and $\text{R}_2 = \text{R}_3 = \text{R}_4 = \text{OH}$ at positions 4, 5 and 8;
 (4) $\text{R}_1 = \text{R}_3 = \text{NH-A-N(O)R'R''}$ at positions 1 and 8 and $\text{R}_2 = \text{R}_4 = \text{OH}$
 20 at positions 4 and 5;
 (5) $\text{R}_1 = \text{R}_2 = \text{NH-A-N(O)R'R''}$ at positions 1 and 4 and $\text{R}_3 = \text{R}_4 = \text{OH}$ at positions 5 and 8; or
 (6) $\text{R}_1 = \text{R}_3 = \text{NH-A-N(O)R'R''}$ at positions 1 and 5 and $\text{R}_2 = \text{R}_4 = \text{OH}$ at positions 4 and 8.
8. A compound according to Claim 6, in which
 (1) $\text{R}_1 = \text{NH-A-N(O)R'R''}$ at position 1, $\text{R}_2 = \text{NH-A-NHR}$ at position 4, and $\text{R}_3 = \text{R}_4 = \text{OH}$ at positions 5 and 8;
 (2) $\text{R}_1 = \text{NH-A-N(O)R'R''}$ at position 1, $\text{R}_2 = \text{OH}$ at position 4, $\text{R}_3 = \text{NH-A-NHR}$ at position 5 and $\text{R}_4 = \text{OH}$ at position 8, or
 30 (3) $\text{R}_1 = \text{NH-A-N(O)R'R''}$ at position 1, $\text{R}_2 = \text{R}_3 = \text{OH}$ at positions 4 and 5 and $\text{R}_4 = \text{NH-A-NHR}$ at position 8.
9. A compound according to Claim 1, in which
 $\text{R}_1 = \text{R}_2 = \text{NH-A-N(O)R'R''}$ at positions 1 and 4 and $\text{R}_3 = \text{R}_4 = \text{OH}$ at positions 5 and 8, or $\text{R}_1 = \text{R}_3 = \text{NH-A-N(O)R'R''}$ at positions 1 and 5
 35 and $\text{R}_2 = \text{R}_4 = \text{OH}$ at positions 4 and 8 with both NH-A-N(O)R'R'' being $\text{NH-(CH}_2)_2\text{N(O)(CH}_3)_2$ or $\text{NH-(CH}_2)_2\text{N(O)(CH}_2\text{CH}_2\text{OH)}_2$.

- 30 -


10. A compound according to Claim 1, in which $R_1 = \text{NH-A-N(O)R'R''}$ at position 1, $R_2 = \text{NH-A-NHR}$ at position 4 and $R_3 = R_4 = \text{OH}$ at positions 5 and 8 or $R_1 = \text{NH-A-N(O)R'R''}$ at position 1, $R_2 = \text{OH}$ at position 4, $R_3 = \text{NH-A-NHR}$ at position 5 and $R_4 = \text{OH}$ at position 8 with NH-A-N(O)R'R'' being $\text{NH-(CH}_2)_2\text{N(O)(CH}_3)_2$ or $\text{NH-(CH}_2)_2\text{N(O)(CH}_2\text{CH}_2\text{OH)}_2$ and NH-A-NHR being $\text{NH-(CH}_2)_2\text{NHCH}_3$ or $\text{NH(CH}_2)_2\text{NHCH}_2\text{CH}_2\text{OH}$.

11. A process for the preparation of a compound of formula (I) as defined in Claim 1 which comprises oxidizing a compound of formula (Ia)



in which R^1 , R^2 , R^3 and R^4 correspond to R_1 , R_2 , R_3 and R_4 , respectively, in the compound (I) but with each of the groups of the type NH-A-N(O)R'R'' in the compound (I) being instead a group NH-A-NR'R'' in the compound (Ia) and one or more groups X, R, R' and R'' in the compound (Ia) optionally instead being in a form convertible to said group or groups present in the compound (I), and where appropriate converting the one or more modified groups X, R, R' and R'' in the compound (Ia) to the form present in the compound (I) and/or forming an acid addition salt with a physiologically acceptable organic or inorganic acid.

12. A compound of formula



- 31 -

in which R_1 , R_2 , R_3 and R_4 are each separately selected from hydrogen, X, NH-A-NHR and NH-A-NR'R" wherein X is hydroxy, halogeno, amino, C₁₋₄ alkoxy or C₂₋₈ alkanoyloxy, A is a C₂₋₄ alkylene group with a chain length between NH and NHR or NR'R" of at least 2 carbon atoms and R, R' and R" are each separately selected from C₁₋₄ alkyl groups and C₂₋₄ hydroxyalkyl and C₂₋₄ dihydroxyalkyl groups in which the carbon atom attached to the nitrogen atom does not carry a hydroxy group and no carbon atom is substituted by two hydroxy groups, or R' and R" together are a C₂₋₆ alkylene group which with the nitrogen atom to which R' and R" are attached forms a heterocyclic group having 3 to 7 atoms in the ring, but with the proviso that at least one of R_1 to R_4 is a group NH-A-NR'R" and at least one other is a different group NH-A-NR'R" or a group NH-A-NHR, the compound optionally being in the form of a physiologically acceptable salt.

13. A compound according to Claim 12, in which R_1 , R_2 , R_3 and R_4 are each separately selected from hydrogen, hydroxy, NH-A-NHR and NH-A-NR'R".

14. A compound according to Claim 12 or 13, in which at least one of R_1 to R_4 is a group NH-A-NR'R" and at least one other is a group NH-A-NHR.

15. A compound according to Claim 14, in which

(1) R_1 = NH-A-NR'R" at position 1, R_2 = NH-A-NHR at position 4, and R_3 = R_4 at positions 5 and 8;

(2) R_1 = NH-A-NR'R" at position 1, R_2 = OH at position 4, R_3 = NH-A-NHR at position 5 and R_4 = OH at position 8, or

(3) R_1 = NH-A-NR'R" at position 1, R_2 = R_3 = OH at positions 4 and 5 and R_4 = NH-A-NHR at position 8.

16. A compound according to any of Claims 12 to 15, in which A is ethylene.

17. A compound according to any of Claims 12 to 15, in which R, R' and R" are each separately selected from CH₃, CH₂CH₃, CH₂CH₂CH₃, CH₂CH₂OH, CH₂CH₂CH₂OH, CH(CH₃)CH₂OH and CH₂CHOHCH₂OH.

- 32 -

18. A pharmaceutical composition comprising a compound of formula (I) as defined in any of Claims 1 to 10 together with a physiologically acceptable diluent or carrier.
19. A compound of formula (I) as defined in any of Claims 1 to 10
05 for use in therapy.
20. A method suitable for aiding regression and palliation of cancer which comprises administering to a patient a therapeutically effective amount of a compound of formula (I) as defined in any of Claims 1 to 10.

1/1




Fig.1a




Fig.1b




Fig.1c




Fig.1d

INTERNATIONAL SEARCH REPORT

International Application No PCT/GB 90/01574

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC ⁵ : C 09 B 1/515		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
IPC ⁵	C 09 B, C 07 C	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	GB, A, 2036554 (ALLIED CHEMICAL) 2 July 1980 see claim 1 --	1
A	US, A, 4138415 (MURDOCK et al.) 6 February 1979 see claim 1 --	1
A	EP, A, 0145226 (WARNER-LAMBERT COMPANY) 19 June 1985 see claim 1 -----	1
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>⁹ Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Z" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
13th December 1990	10. 01. 91	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="font-family: cursive; font-size: 1.2em;">M. Peis</div> <div style="border: 1px solid black; padding: 2px 5px;">M. PEIS</div> </div>	

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

GB 9001574
SA 40975

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 04/01/91. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A- 2036554	02-07-80	GB-A, B 2000029	04-01-79
US-A- 4138415	06-02-79	AT-B- 359484	10-11-80
		AU-A- 3877678	14-02-80
		CA-A- 1099710	21-04-81
		CH-A- 644840	31-08-84
		DE-A- 2835661	01-03-79
		FR-A, B 2400504	16-03-79
		GB-A, B 2004293	28-03-79
		JP-A, B 1063556	09-03-89
		NL-A- 7808475	19-02-79
		SE-B- 445996	04-08-86
		SE-A- 7807987	16-02-79
EP-A- 0145226	19-06-85	US-A- 4626540	02-12-86
		AU-B- 573639	16-06-88
		AU-A- 3513184	16-05-85
		CA-A- 1258856	29-08-89
		DE-A- 3474446	10-11-88
		JP-A- 60136567	20-07-85